BADR UNIVERSITY IN ASSIUT
جـامـعـة بـدر بأسـيـــوط

NEONATAL MANAGEMENT IN VETERINARY MEDICINE

Submitted as Partial Fulfillment of Course Requirements

Submitted by:
Mahmoud Ali El-Sayed Aboulila (Leader)
Diana Sherif
Amna Haggag Mohamed
Huda Abdelshafi
Aya Ahmed El-Sayed
Aya Ali Mohamed
Mariam Mahmoud Ibrahim

Level / Year:
Fourth Year

Course Title:
Internal Diseases

Submitted to:
Prof.\ Nasr El Areef

Faculty:
Faculty of Veterinary Medicine

University:
Badr University in Assuit (BUA)

Table of Contents

Introduction 2

1. Objectives of Neonatal Management 3

2. Physiological Characteristics of Neonates 3

3. Immediate Care at Birth 3

4. Thermoregulation and Environmental Management 4

5. Colostrum Management 4

6. Nutrition of Neonates 5

7. Housing and Hygiene 5

8. Common Neonatal Diseases 5

9. Species-Specific Management 6

10. Monitoring Neonatal Health 6

11. Emergency Neonatal Care 7

12. Prevention Programs 7

13. Economic Importance 7

Conclusion 8

References 9

Introduction

Neonatal management in veterinary medicine refers to the complete care, supervision, and handling of newborn animals during the first hours, days, and weeks of life. This stage is considered one of the most delicate and critical periods in the life of any animal because the newborn must rapidly adapt from the protected intrauterine environment to the external world. During this transition, the neonate becomes responsible for independent breathing, maintaining body temperature, searching for food, and resisting environmental pathogens. Any failure in these early adaptations may lead to weakness, disease, poor growth, or even death.

The neonatal period is associated with a high risk of morbidity and mortality in many animal species. Newborn animals are highly vulnerable to infectious diseases, environmental stress, hypothermia, dehydration, starvation, congenital defects, trauma, and management errors. Their immune systems are immature, their energy reserves are limited, and their ability to regulate body temperature is not fully developed. Therefore, even minor problems that may seem simple in adult animals can become life-threatening in neonates if not recognized and treated quickly.

Proper neonatal management plays a major role in improving survival rate, health status, welfare, and long-term productivity. Early care such as ensuring a clear airway, stimulating breathing, drying the body, disinfecting the umbilical cord, and providing colostrum at the correct time can dramatically increase the chances of survival. In farm animals, successful neonatal care leads to better growth performance, lower treatment costs, improved reproductive efficiency in the future, and greater economic return to the producer. In companion animals such as puppies and kittens, it improves health, vitality, and quality of life.

The neonatal period varies among species, but generally includes the first **2 to 4 weeks after birth**, although some physiological adjustments continue beyond this time. During this period, body systems such as thermoregulation, immunity, digestion, circulation, nervous coordination, and locomotion are still developing. The digestive tract must begin processing milk, the cardiovascular system must adapt to life outside the uterus, and behavioral responses such as standing, suckling, and bonding with the mother must be established rapidly.

Management practices also differ according to species, breed, environment, and production system. For example, piglets require careful temperature support and protection from crushing by the sow, calves need rapid colostrum intake and navel care, foals require close monitoring for standing and nursing behavior, while puppies and kittens need constant warmth and frequent feeding if orphaned. Understanding these species differences is essential for every veterinarian and animal caretaker.

This report discusses the fundamental principles of neonatal management in domestic animals including calves, foals, lambs, kids, puppies, kittens, and piglets. It highlights immediate care after birth, nutritional management, colostrum feeding, environmental control, common neonatal diseases, preventive strategies, and the veterinary role in ensuring healthy development during the earliest and most vulnerable stage of life.

1. Objectives of Neonatal Management

The main goals are:

- Ensure survival of the newborn.
- Promote healthy growth and development.
- Prevent infectious diseases.
- Provide adequate nutrition.
- Maintain body temperature.
- Detect congenital abnormalities early.
- Reduce neonatal mortality and economic losses.

2. Physiological Characteristics of Neonates

Newborn animals differ from adults in several aspects:

a) Poor Thermoregulation

Neonates have:

- Thin skin
- Low body fat reserves
- Large surface area relative to body size

Therefore, they lose heat rapidly and are prone to **hypothermia**.

b) **Immature Immune System:** Most domestic animals are born with little or no circulating antibodies because placental transfer is limited. They depend mainly on **colostrum** for passive immunity.

c) **Limited Energy Reserves:** They possess low glycogen stores and can develop **hypoglycemia** quickly if not fed.

d) **Immature Digestive Function:** Digestive enzymes and rumen development are incomplete (especially in ruminants).

3. Immediate Care at Birth

a) Clearing Airways

Immediately after delivery:

- ✓ Remove fetal membranes from mouth and nostrils.
- ✓ Wipe mucus from nose.
- ✓ Stimulate breathing by rubbing the body with a towel.
- ✓ If necessary, perform gentle suction.

b) Stimulating Respiration

If respiration is weak:

- Rub thorax vigorously.
- Tickle nostrils with straw.
- Provide oxygen if available.

c) **Drying the Neonate:** Drying prevents heat loss and stimulates circulation.

d) Umbilical Cord Care

- Dip umbilical stump in **7% iodine** or chlorhexidine.
- Prevent omphalitis (navel infection).

e) Initial Examination

Check for:

- ✓ Normal breathing
- ✓ Heart rate
- ✓ Reflexes

- ✓ Limb deformities
- ✓ Cleft palate
- ✓ Hernia
- ✓ Weakness

4. Thermoregulation and Environmental Management

Maintaining suitable environmental temperature is essential.

Recommended Temperatures

Species	Ideal Neonatal Temperature
Piglets	30–34°C
Puppies	29–32°C (first week)
Kittens	29–32°C
Lambs/Kids	25–30°C
Calves	15–25°C

⚠ **Methods of Warming**

- Heat lamps
- Incubators
- Warm blankets
- Warm water bottles (carefully wrapped)
- Dry bedding

☒ Avoid overheating and dehydration.

5. Colostrum Management

Importance of Colostrum

Colostrum provides:

- ☑ Immunoglobulins (IgG)
- ☑ Energy
- ☑ Vitamins A, D, E
- ☑ Growth factors
- ☑ Laxative effect (helps pass meconium)

Timing

Absorption of antibodies decreases rapidly after birth:

- ✓ Highest during first **2 hours**
- ✓ Reduced after **12 hours**
- ✓ Gut closure by **24 hours**

Recommended Amount

Calves: 10% of body weight within first 6 hours.

Foals: 1–2 liters within first few hours.

Puppies/Kittens: Nurse within first 12 hours.

Failure of Passive Transfer (FPT): Occurs when inadequate colostrum is consumed.

Consequences:

- Septicemia
- Diarrhea
- Pneumonia
- Death

Diagnosis in foals/calves via serum IgG testing.

6. Nutrition of Neonates

Natural Feeding: Mother's milk is ideal.

Artificial Feeding: Used when:

- ⊠ Maternal death
- ⊠ Rejection
- ⊠ Mastitis
- ⊠ Insufficient milk

Use species-appropriate milk replacer.

Feeding Frequency

Species	Frequency
Puppies/Kittens	Every 2–4 hrs
Piglets	Frequent suckling
Lambs/Kids	4–6 feeds/day
Calves	2–3 feeds/day

Avoid overfeeding to prevent diarrhea.

7. Housing and Hygiene

Good Housing Should Provide:

- Warmth
- Dryness
- Ventilation
- Clean bedding
- Protection from predators/injury

Hygiene Measures

- Disinfect pens regularly
- Clean feeding bottles
- Isolate sick neonates
- Proper manure disposal

Poor hygiene increases risk of neonatal diarrhea and septicemia.

8. Common Neonatal Diseases

a) Neonatal Diarrhea (Scours)

Caused by:

- Escherichia coli infection
- Rotavirus infection
- Salmonellosis
- Nutritional errors

Signs:

- Dehydration
- Weakness
- Sunken eyes

Treatment:

- Oral/IV fluids
- Antibiotics if bacterial
- Warmth

b) **Septicemia:** Systemic bacterial infection due to poor immunity.

Signs:

- Fever or low temperature
- Depression
- Poor suckling

Emergency treatment required.

c) **Pneumonia:** Due to poor ventilation or aspiration.
d) **Navel Ill:** Umbilical infection causing swollen painful navel.
e) **Hypoglycemia:** Common in weak or orphan neonates.

Signs:

- Tremors
- Weakness
- Seizures

9. Species-Specific Management

Calves:

- Stand within 1 hour.
- Nurse within 2 hours.
- Navel dipping.
- Individual clean pen.

Foals:

- Stand in 1 hour.
- Nurse in 2 hours.
- Pass meconium in 3 hours.
- Watch for dummy foal syndrome.

Piglets:

- Prevent crushing by sow.
- Provide creep heat.
- Ensure colostrum intake.
- Iron supplementation at 3–5 days.

Puppies/Kittens:

- Monitor daily weight gain.
- Keep warm.
- Stimulate urination/defecation manually if orphaned.

Lambs/Kids:

- Dry immediately after birth.
- Ensure suckling.
- Prevent starvation-exposure complex.

10. Monitoring Neonatal Health

Daily observation:

- Body temperature
- Weight gain
- Nursing behavior
- Hydration
- Feces consistency
- Activity level

Failure to gain weight is an early warning sign.

11. Emergency Neonatal Care

Indications:

- ✓ Weakness
- ✓ No suck reflexes
- ✓ Hypothermia
- ✓ Dehydration
- ✓ Dyspnea

Immediate Actions:

1. Warm gradually.
2. Correct blood glucose.
3. Give fluids.
4. Provide oxygen.
5. Veterinary examination.

12. Prevention Programs

- ⚠ Vaccinate dams before parturition.
- ⚠ Good maternal nutrition.
- ⚠ Clean maternity area.
- ⚠ Adequate supervision during birth.
- ⚠ Early colostrum administration.
- ⚠ Reduce overcrowding.

13. Economic Importance

Good neonatal management leads to:

- Lower mortality
- Better growth rates
- Reduced treatment costs
- Improved future reproduction
- Higher farm profitability

Conclusion

Neonatal management is a cornerstone of successful veterinary practice and efficient animal production systems. The newborn period represents one of the most sensitive and vulnerable stages of life, during which the animal undergoes major physiological and environmental adjustments in a very short time.

During this stage, even small mistakes in temperature control, feeding management, hygiene, housing conditions, or colostrum intake may rapidly result in weakness, disease, poor growth, or death.

Effective neonatal care requires continuous observation, rapid intervention, and a clear understanding of the special needs of each species. A veterinarian must be able to recognize danger signs early, such as failure to suckle, weakness, dehydration, abnormal body temperature, respiratory distress, or poor weight gain. Early diagnosis and prompt treatment are essential to reduce mortality and prevent long-term complications.

In addition, preventive medicine plays a major role in neonatal success. Good maternal nutrition, vaccination of dams, clean birthing environments, proper assistance during parturition, and immediate postnatal care all contribute significantly to healthier neonates and stronger immunity. These measures are often more effective and economical than treating disease after it occurs.

Proper neonatal management not only saves lives but also improves animal welfare, enhances growth performance, reduces medical costs, and increases future reproductive and productive potential. Healthy neonates are more likely to become strong, productive adults, whether in livestock production systems or companion animal care.

In conclusion, neonatal management should always be considered a top priority in veterinary medicine, because the quality of care provided during the first hours and days of life often determines the future health, performance, and survival of the animal.

References

1. Radostits, O. M., Gay, C. C., Hinchcliff, K. W., & Constable, P. D. (2007). *Veterinary Medicine: A Textbook of the Diseases of Cattle, Horses, Sheep, Pigs and Goats* (10th ed.). Elsevier.
2. Smith, B. P. (2014). *Smith's Large Animal Internal Medicine* (5th ed.). Mosby Elsevier.
3. Peterson, M. E., & Kutzler, M. A. (2011). *Small Animal Pediatrics: The First 12 Months of Life*. Elsevier.
4. Youngquist, R. S., & Threlfall, W. R. (2007). *Current Therapy in Large Animal Theriogenology* (2nd ed.). Saunders Elsevier.
5. Noakes, D. E., Parkinson, T. J., & England, G. C. W. (2018). *Veterinary Reproduction and Obstetrics* (10th ed.). Elsevier.
6. Merck Veterinary Manual. (2025). *Care of newborn animals and neonatal disorders*.
7. World Organisation for Animal Health (WOAH). (2024). *Animal welfare guidelines for neonatal livestock care*.
8. American Veterinary Medical Association (AVMA). (2024). *Guidelines for newborn animal care and welfare*.
9. Food and Agriculture Organization of the United Nations (FAO). (2023). *Livestock production and newborn management practices*.
10. Fubini, S. L., & Ducharme, N. G. (2016). *Farm Animal Surgery* (2nd ed.). Elsevier.
11. Ettinger, S. J., Feldman, E. C., & Côté, E. (2017). *Textbook of Veterinary Internal Medicine* (8th ed.). Elsevier.
12. Pugh, D. G., & Baird, A. N. (2012). *Sheep and Goat Medicine* (2nd ed.). Elsevier.
13. Taylor, D. J. (2013). *Pig Diseases* (9th ed.). 5M Publishing.
14. Kahn, C. M., & Line, S. (Eds.). (2010). *The Merck/Merial Manual for Pet Health*. Merck.
15. Madigan, J. E., & McCue, P. M. (2015). *Equine Neonatal Medicine and Surgery*. Wiley-Blackwell.